

TIPS

TO HELP NAVIGATE FINANCIAL ASSISTANCE

INDICATION

AUBAGIO® (teriflunomide) is a prescription medicine used to treat relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.

IMPORTANT SAFETY INFORMATION

Do not take AUBAGIO if you have severe liver problems, are pregnant or of childbearing potential and not using effective birth control, have had an allergic reaction to AUBAGIO or leflunomide, or are taking a medicine called leflunomide for rheumatoid arthritis.

Please see Important Safety Information and full [Prescribing Information](#), including boxed **WARNING** and [Medication Guide](#).



Once-daily

AUBAGIO[®]
(teriflunomide) 14mg
Tablets

AUBAGIO is available in 14 mg and 7 mg tablets.

GETTING PERSONALIZED SUPPORT



When it comes to paying for your disease-modifying treatment, dealing with insurance and understanding your coverage can be complicated.¹

Keep reading to learn more about financial assistance and support programs for AUBAGIO.



MS One to One[®] helped navigate my insurance approval. They have a whole department for that! No extra work for me."

- Heather

On AUBAGIO 14 mg
Individual experiences may vary.



FINANCIAL ASSISTANCE FOR AUBAGIO



ENROLL IN THE \$0 CO-PAY PROGRAM

The AUBAGIO Co-Pay Program offers assistance for those with commercial insurance. It assists with co-payments and co-insurance amounts for AUBAGIO, regardless of your financial status. Once enrolled in the Program, you'll have \$0 co-payments for AUBAGIO.*



SPEAK TO AN MS ONE TO ONE NURSE

For people taking AUBAGIO, *MS One to One* offers assistance navigating insurance, including benefits verification and financial assistance for eligible patients. Your Nurse can also help you connect with outside financial assistance programs.

Please see Important Safety Information and full [Prescribing Information](#), including boxed **WARNING** and [Medication Guide](#).

*Out-of-pocket costs related to medical appointments, evaluations, testing, or other related services are not covered by the AUBAGIO Co-Pay Program. The Co-Pay Program is not available for prescriptions purchased under Medicare, Medicaid, TRICARE, or other federal- and state-funded programs.



ONE START[®] HELPS SIMPLIFY STARTING AUBAGIO

With this program, eligible commercially insured patients can begin treatment within 5 business days of being prescribed AUBAGIO. If needed, you can receive AUBAGIO at no cost to you for up to 1 year while your benefits are being verified.

The *One Start* Program is not available for prescriptions purchased under Medicare, Medicaid, TRICARE, or other federal- and state-funded programs.

MORE FINANCIAL RESOURCES

If you don't have insurance or are unable to pay for AUBAGIO, ask an *MS One to One*[®] Nurse about the Sanofi Genzyme Patient Assistance Program, or check with your local patient advocacy group about other support options.

Please see Important Safety Information and full [Prescribing Information](#), including boxed **WARNING** and [Medication Guide](#).

THANK YOU FOR CHOOSING AUBAGIO.



A Community Sponsored by
AUBAGIO[®] (teriflunomide)

JOIN THE COMMUNITY

Common Threads offers personal stories, articles, and videos for healthy living while taking AUBAGIO. Get started at aubagiocommunity.com



GIVE US A CALL

MS One to One[®] Nurses are available any time, day or night. Have a question? We have answers. Call us at **1-855-676-6326**



VISIT US ONLINE

Check out aubagio.com for special events, patient stories, and more information about AUBAGIO.

AUBAGIO[®]
(teriflunomide) 14mg tablets

**WE'RE HERE
TO HELP.**

AUBAGIO[®]
(teriflunomide) 14mg
tablets



Call ***MS One to One***[®] at
1-855-676-6326

REFERENCE:

1. Bourdette DN, Hartung DM, Whitham RH. Practices of US health insurance companies concerning MS therapies interfere with shared decision-making and harm patients. *Neurol Clin Pract.* 2016;6(2):177-182.

Please see Important Safety Information and full [Prescribing Information](#), including boxed **WARNING** and [Medication Guide](#).

INDICATION

AUBAGIO® (teriflunomide) is a prescription medicine used to treat relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.

IMPORTANT SAFETY INFORMATION

DO NOT TAKE AUBAGIO IF YOU:

- **Have severe liver problems. AUBAGIO may cause serious liver problems**, including liver failure that can be life-threatening and may require a liver transplant. Your risk may be higher if you already have liver problems or take other medicines that affect your liver. Your healthcare provider should do blood tests to check your liver within 6 months before you start AUBAGIO and monthly for 6 months after starting AUBAGIO. Tell your healthcare provider right away if you develop any of these symptoms of liver problems: nausea, vomiting, stomach pain, loss of appetite, tiredness, yellowing of your skin or whites of your eyes, or dark urine.
- **Are pregnant. AUBAGIO may harm an unborn baby.** You should have a pregnancy test before starting AUBAGIO. After stopping AUBAGIO, continue to use effective birth control until you have made sure your blood levels of AUBAGIO are lowered. If you become pregnant while taking AUBAGIO or within 2 years after stopping, tell your healthcare provider right away and enroll in the AUBAGIO Pregnancy Registry at 1-800-745-4447, option 2.
- **Are of childbearing potential and not using effective birth control. It is not known if AUBAGIO passes into breast milk.** Your healthcare provider can help you decide if you should take AUBAGIO or breastfeed — you should not do both at the same time.
If you are a man whose partner plans to become pregnant, you should stop taking AUBAGIO and talk with your healthcare provider about reducing the levels of AUBAGIO in your blood. If your partner does not plan to become pregnant, use effective birth control while taking AUBAGIO.
- **Have had an allergic reaction to AUBAGIO or a medicine called leflunomide.**
- **Take a medicine called leflunomide for rheumatoid arthritis.**



©2021 Genzyme Corporation. All rights reserved.

AUBAGIO, *MS One to One*, Sanofi and Genzyme registered in U.S. Patent and Trademark Office.

MAT-US-2012825-v3.0-05/2021

Please see Important Safety Information and full [Prescribing Information](#), including boxed **WARNING** and [Medication Guide](#).

AUBAGIO may stay in your blood for up to 2 years after you stop taking it. Your healthcare provider can prescribe a medicine that can remove AUBAGIO from your blood quickly.

Before taking AUBAGIO, talk with your healthcare provider if you have: liver or kidney problems; a fever or infection, or if you are unable to fight infections; numbness or tingling in your hands or feet that is different from your MS symptoms; diabetes; serious skin problems when taking other medicines; breathing problems; or high blood pressure. Your healthcare provider will check your blood cell count and TB test before you start AUBAGIO. Talk with your healthcare provider if you take or are planning to take other medicines (especially medicines for treating cancer or controlling your immune system), over-the-counter medicines, vitamins or herbal supplements.

AUBAGIO may cause serious side effects. Tell your doctor if you have any of the following:

- **decreases in white blood cell count** — this may cause you to have more infections. Symptoms include fever, tiredness, body aches, chills, nausea, or vomiting. Patients with low white blood cell count should not receive certain vaccinations during AUBAGIO treatment and 6 months after.
- **allergic reactions** such as difficulty breathing, itching, or swelling on any part of your body including lips, eyes, throat, or tongue. Stop taking AUBAGIO and call your doctor right away.
- **serious skin reactions** that may lead to death. Stop taking AUBAGIO if you have rash or redness and peeling, mouth sores or blisters.
- **other allergic reactions that may affect different parts of the body.** If you have a fever or rash in combination with severe muscle pain, swollen lymph glands, swelling of your face, unusual bruising or bleeding, weakness or tiredness, or yellowing of your skin or the whites of your eyes, stop taking AUBAGIO and call your doctor right away.
- **numbness or tingling in your hands or feet that is different from your MS symptoms**
- **high blood pressure**
- **breathing problems (new or worsening)** — these may be serious and lead to death

The most common side effects when taking AUBAGIO include: headache; diarrhea; nausea; hair thinning or loss; and abnormal liver test results. These are not all the side effects of AUBAGIO. Tell your healthcare provider about any side effect that bothers you.

Consult your healthcare provider if you have questions about your health or any medications you may be taking, including AUBAGIO.

